

Low latitude ionospheric turbulence observed by Aureol 3 and possible association with seismicity

Hobara et al., 2005 in Ann. Geophys. (In press)


Hobara. Y¹, M. Parrot², F. Lefeuvre² and O.A. Molchanov³

1. Swedish Institute of Space Physics (IRF), Kiruna Sweden
2. LPCE/CNRS, Orléans, France
3. Institute of the Earth Physics, Moscow, Russia.

E-mail : yasu@irf.se

Introduction

Characteristic scale of IT



OBJECTIVES

- IT and electric field turbulence in low ~ middle latitudes.
- Effect to IT from energy sources from the earth (e.g. seismic activities)
- Lithosphere-atmosphere-ionosphere coupling

PREVIOUS WORKS

- Molchanov et al., 2002: 2004 IK24, Cosmos-900
- Cerisier et al., 1985. Aureol 3, High lat IT & E.
- Parrot, 1991, Aureol 3, VLF emissions.
- Galperin et al., 1996. Aureol 3, Ground explosion.

Contents

- Introduction
- Data used
- Results from waveform data (case study)
- Results from filter bank data (statistical study)
- Conclusion

Data from Aureol-3

- Perigee 400km, Apogee 2000km
Inclination 82.5deg, Period 109.5min
- Launch: Sep 1981 Termin: Jan 1986


Data used for current analysis

Electric Field

- Electric field EH (Y -component)
- Waveform(WF): $6\text{Hz} < f < 1500\text{Hz}$
- Filter bank (FBE): $10\text{Hz} < f < 1000 \text{ Hz}$
- Time resolution: 0.01 to 2.56 sec (ZAP1 to ZAP4)

Electron Density



- ISOPROBE (High resolution): $6\text{Hz} < f < 1\text{kHz}$



Waveform and FB


	Waveforms (High time resolution)	Filter banks
Orbit	Near TM station	Entire earth
Component	EH and Ne	EH
Frequency resolution	Arbitrary ($f_s=5\text{kHz}$)	6 channels $(10\text{Hz} < f < 1000\text{Hz})$
	Case Study	Statistical Study

Ne near the Equator (WF)



PSD (WF)

1982/6/8/0:58:10.043 (UT) Mlat=7.81 deg, Alt=437.0 km




Power spectra of (a) electron density and (b) electric field turbulence inside equatorial spread F (ESF)

$\alpha_n \sim \alpha_E$: RT instability regime, $P \propto f^{-\alpha}$ ($6\text{Hz} < f < 100\text{Hz}$)


Spectral index (6Hz<f<100Hz)

$\delta N_e / N_e$
 δE



Spectral index (30Hz<f<300Hz)

$\delta N_e / N_e$
 δE



Statistical study (FB)

Advantage: Global orbital coverage

Data selection

Orbit

- Altitude < 900 km
- - 45 deg < Magnetic latitude < 45 deg

Electric Field Power ($10 \text{ Hz} < f < 100 \text{ Hz}$)

- **Burst event** (significant local enhancement of E-power)
- **Burst intensity** and corresponding **spectral index**


Seismic events

- NGDC seismic catalogue (1964-1995)
- Extract events : $M_s > 5.5$, Depth < 100 km


Burst event (FB) -1

Seismic events:

- $M > 5.5$
- $\Delta \text{Magnetic long} < 15 \text{deg}$
- $\Delta \text{Day} < 5 \text{days}$




1982/03/12 Orbit 2




Mlat [deg]	-15.09	0.06	19.95	40.95
Glong[deg]	148.98	149.88	151.43	154.52
Altitude [km]	879.06	717.04	544.85	433.11
Time [UT]	09:20:24	09:24:19	09:29:24	09:34:34

Burst event (FB) -2




Number of H-orbits:

- Total : 864
- Low alt bursts : 96
- Bursts with EQ: 32




Lat-long distribution of the burst



Spectral index (FB)

*Error bars: 95 % confidence interval (Gaussian distribution)


No EQs
With EQs



Electric Field Power (FB)

*Error bars: 95 % confidence interval (Gaussian distribution)

No EQs
With EQs





The statistical test

The probability to have different parent distributions between seismic and non seismic periods. (Gaussian distribution assumed)

	Sp. and Au.	Su. and Wi.	Mo. and Ev.	Day	Night	All
Spectral index	0.55	0.67	0.55	0.05	0.5	0.094
Electric field power	0.7	0.82	0.61	0.38	0.45	0.93

Probability Distributions

Solid: ALL
Dotted: EQ
Dash dotted: NO EQ



Conclusion

- Spectral indices for density and electric field turbulence in the IT (ESF) show similar value around 2 for different days (case study)
- 96 events of electric field enhancement (burst events) connected to the IT in low-mid latitudes are analyzed.
- Spectral indices for seismic and non-seismic events have no remarkable statistical difference.
- Mean E-field burst power between seismic and non-seismic burst events has a statistical difference with 93 % confidence.
- AGW and/or atmospheric turbulence due to the seismic activity may intensify the IT.